

# RUMEN ACIDOSIS IN DAIRY CATTLE: BUNK MANAGEMENT CONSIDERATIONS

**R. D. Shaver, Ph.D.**

**Professor and Extension Dairy Nutritionist<sup>1</sup>**  
**Department of Dairy Science**  
**College of Agricultural and Life Sciences**  
**University of Wisconsin – Madison**  
**University of Wisconsin – Extension**

## **Introduction**

Laminitis has been linked to lactic acidosis (16). Acidosis is caused by the consumption of high amounts of ruminally-available carbohydrate, low amounts of effective fiber, or both (16). Nordlund (18) and Oetzel (19) reported that sub-acute rumen acidosis (SARA) is a prevalent problem for commercial dairies.

Despite diet formulation for chemical (15) and physically-effective fiber (12) minimums, non-structural carbohydrate (NSC) and starch maximums (16), and the use of total mixed rations (TMR), some degree of SARA may be inevitable in high-producing dairy herds because total chewing and rumination times per kg of rumen-fermentable organic matter (RFOM) intake declines as RFOM intake increases (Figures 1 and 2). Using RFOM intake to predict fermentation acid production and total chewing time to predict salivary buffer flow, the widening difference between acid production and buffer flow as RFOM intake increases is presented in Figure 3. Increased intake of RFOM as milk production increases is a normal consequence of high milk production, because of increases in DM intake and the feeding of higher-concentrate diets to increase dietary energy density (15).

Increases in chewing duration (min/d) at increasing levels of milk production were not proportional to the increase in DM intake, and total chewing and rumination times per kg of DM intake declined as milk production and DM intake increased (5). Total chewing ( $r = -0.77$ ;  $P < 0.01$ ) and rumination ( $r = -0.68$ ;  $P < 0.05$ ) times per kg of DM intake were negatively correlated with milk production (5). These observations explain the relationships presented in Figures 1 – 3, which highlight the difficulty that we face in avoiding SARA in high-producing dairy herds. The foregoing discussion implicates bunk management as a risk factor for SARA in high-producing dairy herds, because margins for error in feeding programs are small.

Shaver (20) discussed feed delivery and bunk management aspects of acidosis and laminitis in dairy herds. Included in the myriad of errors in feed delivery and bunk management that can occur on commercial dairies were:

- errors in feed sampling and analyses,
- errors in ingredient dry matter (DM) adjustments,
- failure to evaluate forage and TMR particle size,

---

<sup>1</sup> Contact at: Room 266 Animal Sciences Building, 1675 Observatory Drive, University of Wisconsin, Madison, WI 53706, (608)263-3491, FAX (608)263-9412, Email: rdshaver@facstaff.wisc.edu

- failure to evaluate grain moisture content and degree of processing,
- errors in ingredient feeding rates,
- mixing errors including over-mixing that causes particle size reduction,
- poor feeding and management of transition cows,
- feed sorting, and
- practices that promote consumption of fewer and larger meals more quickly.

The latter two areas will be discussed in this paper. It should be noted that bunk management practices that cause an increased incidence of SARA may or may not result in an increased incidence of laminitis. Nordlund (17) reported on the diagnosis of SARA without a high corresponding incidence of laminitis in three grazing herds, and attributed this to the fact that the cows were on dirt rather than concrete. Comparing these findings to diagnostic work-ups done in confined herds, Nordlund (17) suggested that the degree of SARA needed to cause laminitis is greater for cows on dirt than for cows with significant exposure to concrete. Colam-Ainsworth et al. (4) reported increased laminitis in cows that spent more time standing on concrete rather than lying in stalls. Mishra et al. (14) reported lower ruminal pH for dairy cows in hot-humid (29.4° C and 85% relative humidity) than cool (18.3° C and 50% relative humidity) environments when fed either high-roughage (35% grain; pH of 6.1 vs. 6.4) or high-grain (65% grain; pH of 5.6 vs. 6.1) diets, possibly because of decreased rumination activity, increased slug feeding, and (or) increased feed sorting associated with the hot-humid environmental conditions. Cow comfort, her environment, and the formulated diet need to be evaluated in conjunction with bunk management practices when investigating laminitis problem herds.

### **Feed Sorting**

Leonardi and Armentano (8) and Martin (10, 11) observed extensive TMR sorting in the feed bunk in university and on-farm trials, respectively. Data on particle size of TMR and orts and DM intake indicated that cows sorted against the coarse particles (8). This sorting against the coarse particles was more evident for the TMR containing 40% compared to 20% alfalfa hay (DM basis), and the variation in sorting among cows was large (8).

Martin (10, 11) determined the particle size of TMR and bunk mix at 6-h intervals post-feeding on a commercial dairy. The percentages on the top screen of the Penn State shaker box (7) for TMR and bunk mix at 6-, 12-, 18-, and 24-hours post-feeding were 9.3, 13.7, 21.5, 27.5, and 58.7%, respectively. Cows sorted against the coarse particles. From a projection of the coarse particle intake at each time period, it appeared that intake of coarse particles was less than predicted during hours 0 – 12 post-feeding and more than predicted during hours 13 – 24 post-feeding.

Leonardi et al. (9) reported that feeding oat silage with increasing particle size increased sorting against coarse TMR particles. Burato et al. (3) reported that cows with the highest pre-trial milk yield had increased sorting against the coarse TMR particle fraction. This response in feeding behavior reduced the original difference in particle size between their long- and short- chopped alfalfa hay diets resulting in no effect of diet particle size on feed intake or milk production.

Factors that may make TMR prone to sorting include: DM content and particle size of forage and mix, variation in bulk density of feed ingredients, large pieces of cobs and husks in the corn silage, amount and quality of hay added to mix, improper sequencing of ingredients into the mixer, frequency of feeding and push-up, availability of bunk space, and bunk access time. An on-farm evaluation of sorting should include particle size determination (7) of TMR, bunk mix, and refusals.

If sorting is determined to be a problem, then one or more of the following options may need to be considered: feeding smaller amounts of TMR more frequently, adding less hay to the mix, processing hay finer, using higher quality hay, using hay that is more pliable, processing corn silage, addition of water to dry TMR, and addition of a liquid feed supplement (LFS) to TMR.

Carver (personal comm.; Quality Liquid Feeds, Dodgeville, WI) determined the particle size of TMR, bunk mix, and refusals (Lammers et al., 1996) on commercial dairies. Data for TMR and refusals are presented in Table 1. There was sorting against the coarse particle fraction for TMR with no LFS added, as shown by the increased percentages retained on the top screen of the Penn State shaker box for refusals. Addition of LFS to TMR appeared to alleviate the problem of sorting against the coarse particle fraction, as shown by the lack of increase in percentages retained on the top screen of the Penn State shaker box for refusals. It should be noted that these would not be considered to be dry TMR, as they averaged about 55% DM across the farms with a range of 51 to 58% DM. A 2 lb. LFS/cow/day TMR inclusion appeared to be more effective than a 1 lb. LFS/cow/day TMR inclusion. Adding LFS as the final ingredient in TMR was more effective for reducing sorting than adding LFS to forage, which was more effective than adding LFS to concentrates. It should be noted that adding LFS to TMR to prevent sorting also adds rumen-fermentable carbohydrate to the diet through molasses and (or) whey. This needs to be evaluated closely relative to the diet that the LFS is being added to so as to not cause, rather than ameliorate, an acidosis problem.

## **Meal Patterns**

Ruminal pH declines following meals with the rate of pH decline increasing as meal size increases and as dietary neutral detergent fiber (NDF) concentration decreases (2). Bunk management practices that cause cows to eat fewer and larger meals more quickly may be associated with an increased incidence of SARA and laminitis. Factors that can cause slug feeding of the TMR include:

- limited bunk space,
- limited feed access time,
- restricted feeding versus feeding for 5% to 10% refusal,
- inconsistent feeding schedule,
- infrequent TMR push up,
- bunk competition, and
- heat stress.

The combination of limited bunk space (< .45 m per cow) and feed-access time (< 16 to 20 hours per day) is worse than either situation alone. The use of lock-ups in situations of limited bunk space and feed-access time exacerbates the problem, because each lock up and the cow in it takes up .61 m of bunk space. When overcrowding of free stalls coincides with limited bunk space, as

is often the case, the potential for laminitis is greater because cows may spend more time standing on concrete rather than lying in stalls (4).

Feeding the TMR in a drive-by bunk at 10 cm above the cow alley rather than in an elevated bunk increases salivary flow and reduces sorting (1), which may help reduce SARA. Milton (13) reported that feedlot cattle fed to a clean bunk had reduced frequency of meals (4.5 versus 8.2 meals per day) and greater average meal size (3.5 versus 1.6 kg per meal) than cattle fed ad libitum. Milton (13) also reported that deviations of 2 to 4 hours from a normal feeding schedule greatly increased the risk of acidosis in feedlot cattle. Frequent TMR push-up may encourage cows to come to the bunk for more frequent consumption of small meals. First lactation heifers fed in a separate group spent 10% to 15% more time eating and consumed 0.5 to 2.0 more meals per day than herd-mates grouped with mature cows (6).

## **Conclusions**

Some degree of SARA may be inevitable in high-producing dairy herds, because total chewing and rumination times per kg of RFOM intake declines as RFOM intake increases. Bunk management is a risk factor for SARA, and a myriad of errors in feed delivery and bunk management can occur on commercial dairies. Bunk management practices that promote feed sorting and slug feeding must be controlled to minimize the incidence of SARA. Cow comfort, her environment, and the formulated diet need to be evaluated in conjunction with bunk management practices when investigating laminitis problem herds.

## **References**


1. Albright, J. L. 1993. Feeding behavior of dairy cattle. *J. Dairy Sci.* 76:485-498.
2. Allen, M. S. 1997. Relationship between fermentation acid production in the rumen and the requirement for physically effective fiber. *J. Dairy Sci.* 80:1447-1462.
3. Burato, G. M., J. A. Voelker, and M. S. Allen. 2001. Effects of pretrial milk yield on feed intake, production, and feeding behavior responses to forage particle size by lactating cows. *J. Dairy Sci.* 84(Suppl.1):(abstr.)
4. Colam-Ainsworth, P., G. A. Lunn, R. C. Thomas, and R. G. Eddy. 1989. Behavior of cows in cubicles and its possible relationship with laminitis in replacement dairy heifers. *Vet. Rec.* 125:573-575.
5. Dado, R. G., and M. S. Allen. 1994. Variation in and relationships among feeding, chewing, and drinking variables for lactating dairy cows. *J. Dairy Sci.* 77:132-144.
6. Krohn, C. C., and S. P. Kongaard. 1979. Effects of isolating first-lactation cows from older cows. *Livestock Prod. Sci.* 6:137-146.
7. Lammers, B. P., D. R. Buckmaster, and A. J. Heinrichs. 1996. A simple method for the analysis of particle sizes of forage and total mixed rations. *J. Dairy Sci.* 79:922-928.

8. Leonardi, C., and L. E. Armentano. 2000. Effect of particle size, quality and quantity of alfalfa hay, and cow on selective consumption by dairy cattle. *J. Dairy Sci.* 83(Suppl.1):272 (abstr.)
9. Leonardi, C., L. E. Armentano, and K. J. Shinnars. 2001. Effect of different particle size distribution of oat silage on feeding behavior and productive performance of dairy cattle. *J. Dairy Sci.* 84(Suppl.1):(abstr.)
10. Martin, R. 2000. Evaluating TMR particle distribution: A series of on-farm case studies. Pages 75-88 in Proc. 4-State Prof. Dairy Mgmt. Seminar. Dubuque, IA. MWPS-4SD8. MidWest Plan Serv. Ames, IA.
11. Martin, R. 1999. TMR particle distribution analysis at six hour time intervals. Pages 7-16 in Proc. UW Arlington Dairy Day. Arlington, WI. Dairy Sci. Dept., Univ. of WI-Madison.
12. Mertens, D. R. 1997. Creating a system for meeting the fiber requirements of dairy cows. *J. Dairy Sci.* 80:1463-1481.
13. Milton, T. 1998. Feed bunk and feed ingredient management: Perspectives from the beef feedlot industry. Pages 222-229 in Proc. Dairy Feeding Systems Mgmt., Components, and Nutr. Conf. Camp Hill, PA. NRAES-116. Nat. Res., Ag., and Eng. Serv., Coop. Ext., Ithaca, NY.
14. Mishra, M., F. A. Martz, R. W. Stanley, H. D. Johnson, J. R. Campbell, and E. Hilderbrand. 1970. Effect of diet and ambient temperature-humidity on ruminal pH, oxidation reduction potential, ammonia and lactic acid in lactating cows. *J. Animal Sci.* 30:1023-1028.
15. National Research Council. 2001. Nutrient Requirements of Dairy Cattle. 7<sup>th</sup> rev. ed. Natl. Acad. Sci., Washington, DC.
16. Nocek, J. E. 1997. Bovine acidosis: Implications on laminitis. *J. Dairy Sci.* 80:1005-1028.
17. Nordlund, K. 2000. Sore feet, sour rumens, clinical quandaries. Pages 58-64 in Proc. 33<sup>rd</sup> Annu. Conv. Of Amer. Assoc. Bov. Pract. Rapid City, SD.
18. Nordlund, K. 1995. Questions and answers regarding rumenocentesis and the diagnosis of herd-based subacute rumen acidosis. Pages 88-101 in Proc. 4-State Applied Nutr. and Mgmt. Conf. La Crosse, WI. Univ. IA, IL, MN, and WI Coop Ext Serv.
19. Oetzel, G. R. 1997. Using rumenocentesis to diagnose subacute ruminal acidosis in dairy herds. Pages 8-11 in Proc. 4-State Applied Nutr. and Mgmt. Conf. La Crosse, WI. Univ. IA, IL, MN, and WI Coop. Ext. Serv.
20. Shaver, R. D. 2000. Feed delivery and bunk management aspects of laminitis in dairy herds fed total mixed rations. Pages 70-77 in Proc. III Int. Conf. on Bovine Lameness. Parma, Italy.


21. Yang, W. Z., K. A. Beauchemin., and L. M. Rode. 2000, Effects of barley grain processing on extent of digestion and milk production of lactating cows. *Journal of Dairy Sci.* 83: 554-568.

**Table 1.** Percentages of TMR and refusals on top screen of Penn State shaker box from five dairy farms (L. Carver, personal comm., QLF).


Dairy	No LFS	No LFS	LFS	LFS
	<u>TMR</u>	<u>Refusals</u>	<u>TMR</u>	<u>Refusals</u>
	<u>%</u>	<u>%</u>	<u>%</u>	<u>%</u>
A	9.6	26.7	15.8	8.5
B	3.1	16.7	7.8	8.1
C	16.5	53.5	34.7	27.2
D	33.5	52.8	26.7	21.8
E	25.0	55.0	33.0	25.0



**Figure 1.** Relationship between total chewing time per kilogram of rumen-fermentable organic matter intake and intake of rumen fermentable organic matter.



**Figure 2.** Relationship between rumination time per kilogram of rumen-fermentable organic matter intake and intake of rumen fermentable organic matter.



**Figure 3.** Calculated fermentation acid production and salivary buffer flow at varying intakes of rumen-fermentable organic matter.